

Removal of Biological Contaminants

Technical Field

The present invention relates to methods for the removal of biological contaminants, particularly removal of biological contaminants from biological preparations.

Background Art

The modern biotechnology industry is faced with a number of problems especially concerning the processing of complex biological solutions which ordinarily include proteins, nucleic acid molecules and complex sugars and which are contaminated with unwanted biological materials. Contaminants include microorganisms such as bacteria and viruses or biomolecules derived from microorganisms or the processing procedure. The demand is, therefore, for a high purity, scalable separation, which can be confidently used both in product development and production, which in one step will both purify macromolecules and separate these biological contaminants.

Viruses are some of the smallest non-cellular organisms known. These simple parasites are composed of nucleic acid and a protein coat. Viruses are typically very small and range in size from 1.5×10^{-8} m to 5.0×10^{-5} m. Viruses depend on the host cells that they infect to reproduce by inserting their genetic material into the host. Often literally taking over the host's function. An infected cell produces more viral protein and genetic material, often instead of its usual products. Some viruses may remain dormant inside host cells. However, when a dormant virus is stimulated, it can enter the lytic phase where new viruses are formed. Self-assemble occurs and burst out of the host cell results in killing the cell and releasing new viruses to infect other cells. Viruses cause a number of diseases in humans including smallpox, the common cold, chicken pox, influenza, shingles, herpes, polio, rabies. Ebola, hanta fever, and AIDS. Some types of cancer have been linked to viruses.

Pyrogens are agents which induce fever. Bacteria are a common source for the production of endotoxins which are pyrogenic agents. Furthermore, another detrimental effect of endotoxins is their known adjuvant effect which could potentially intensify immune responses against therapeutic drugs. The endotoxin limit set by the Food and Drug Administration (FDA) guidelines for most pharmaceutical products is for a single dose 0.5ng endotoxin per

kilogram body weight or 25ng endotoxin/dose for a 50kg adult. Due to their size and charge heterogeneity, separation of endotoxins from proteins in solution can often be difficult.

Endotoxin inactivation by chemical methods are unsuitable because they are stable under extremes of temperature and pH which would destroy the proteins. Furthermore, due to their amphipathic nature, endotoxins tend to adhere to proteins in a fashion similar to detergents. In such cases, endotoxin activity often clusters with the protein when chromatographic procedures such as ion exchange chromatography or gel filtration are employed.

Presently, the purification of biomolecules is sometimes a long and cumbersome process especially when purifying blood proteins. The process is made all the more complex by the additional step of ensuring the product is “bug” free. The costs associated with this task is large and further escalates the purification costs in total. The Gradiflow technology rapidly purifies target proteins with high yield. For example, a proteins like fibrinogen (a clotting protein) can be separated in three hours using the Gradiflow while the present industrial separation is 3 days. Certain monoclonal antibodies can be purified in 35 minutes compared to present industrial methods which take 35 hours.

The membrane configuration in the Gradiflow enables the system to be configured so that the purification procedure can also include the separation of bacteria viruses and vectors. It has now been found by the present inventors that appropriate membranes can be used and the cartridge housing the membrane configured to include separate chambers for the isolated bacteria and viruses.

The Gradiflow Technology

Gradiflow is a unique preparative electrophoresis technology for macromolecule separation which utilises tangential flow across a polyacrylamide membrane when a charge is applied across the membrane (AU 601040). The general design of the Gradiflow system facilitates the purification of proteins and other macromolecules under near native conditions. This results in higher yields and excellent recovery.

In essence the Gradiflow technology is bundled into a cartridge comprising of three membranes housed in a system of specially engineered grids and gaskets which allow separation of macromolecules by charge and/or molecular weight. The system can also concentrate and desalt/dialyse at the same time. The multimodal nature of the system allows this technology to be

used in a number of other areas especially in the production of biological components for medical use. The structure of the membranes may be configured so that bacteria and viruses can be separated at the point of separation - a task which is not currently available in the biotechnology industry and adds to the cost of production through time delays and also because of the complexity of the task.

Disclosure of invention

In a first aspect, the present invention consists in a method of removing a biological contaminant from a mixture containing a biomolecule and the biological contaminant, the method comprising:

- (a) placing the biomolecule and contaminant mixture in a first solvent stream, the first solvent stream being separated from a second solvent stream by an electrophoretic membrane;
- (b) selecting a buffer for the first solvent stream having a required pH;
- (c) applying an electric potential between the two solvent streams causing movement of the biomolecule through the membrane into the second solvent stream while the biological contaminant is substantially retained in the first sample stream, or if entering the membrane, being substantially prevented from entering the second solvent stream;
- (d) optionally, periodically stopping and reversing the electric potential to cause movement of any biological contaminants having entered the membrane to move back into the first solvent stream. wherein substantially not causing any biomolecules that have entered the second solvent stream to re-enter first solvent stream: and
- (e) maintaining step (c), and optional step (d) if used, until the second solvent stream contains the desired purity of biomolecule.

In a second aspect, the present invention consists in a method of removing a biological contaminant from a mixture containing a biomolecule and the biological contaminant, the method comprising:

- (a) placing the biomolecule and contaminant mixture in a first solvent stream, the first solvent stream being separated from a second solvent stream by an electrophoretic membrane;
- (b) selecting a buffer for the first solvent stream having a required pH;

(c) applying an electric potential between the two solvent streams causing movement of the biological contaminant through the membrane into the second solvent stream while the biomolecule is substantially retained in the first sample stream, or if entering the membrane, being substantially prevented from entering the second solvent stream;

5 (d) optionally, periodically stopping and reversing the electric potential to cause movement of any biomolecule having entered the membrane to move back into the first solvent stream, wherein substantially not causing any biological contaminants that have entered the second solvent stream to reenter first solvent stream; and

(e) maintaining step (c), and optional step (d) if used, until the first solvent stream
10 contains the desired purity of biomolecule.

In the first and second aspects of the present invention, preferably the biomolecule is selected from the group consisting of blood protein, immunoglobulin, and recombinant protein.

The biological contaminant can be a virus, bacterium, prion or an unwanted biomolecule such as lipopolysaccharide, toxin or endotoxin.

Preferably, the biological contaminant is collected or removed from the first stream.

Preferably, the buffer for the first solvent stream has a pH lower than the isoelectric point of biomolecule to be separated.

In a further preferred embodiment of the first aspect of the present invention, the electrophoretic membrane has a molecular mass cut-off close to the apparent molecular mass of biomolecule. It will be appreciated, however, that the membrane may have any required
20 molecular mass cut-off depending on the application. Usually, the electrophoretic membrane has a molecular mass cut-off of between about 3 and 1000kDa. A number of different membranes may also be used in a desired or useful configuration.

The electric potential applied during the method is selected to ensure the required
25 movement of the biomolecule, or contaminant if appropriate, through the membrane. An electric potential of up to about 300 volts has been found to be suitable. It will be appreciated, however, that greater or lower voltages may be used.

The benefits of the method according to the first aspect of the present invention are the possibility of scale-up, and the removal of biological contaminants present in the starting
30 material without adversely altering the properties of the purified biomolecule.

In a third aspect, the present invention consists in use of Gradiflow in the purification or

separation of biomolecule from a biological contaminant.

In a fourth aspect, the present invention consists in biomolecule substantially free from biological contaminants purified by the method according to the first aspect of the present invention.

5 In a fifth aspect, the present invention consists in use of biomolecule according to the third aspect of the present invention in medical and veterinary applications.

In a sixth aspect, the present invention consists in a substantially isolated biomolecule substantially free from biological contaminants.

10 Throughout this specification, unless the context requires otherwise, the word “comprise”, or variations such as “comprises” or “comprising”, will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

In order that the present invention may be more clearly understood a preferred forms will be described with reference to the accompanying drawings.

Brief Description of Drawings

Figure 1. Samples from up and downstream were taken at time intervals (x-axis) during the isolation of albumin from plasma. Albumin was measured in the samples by mixing with BCG reagent and reading the absorbance of 630nm. The concentration of albumin in each sample was calculated from the standard curve, and multiplied by the volume of the up-or downstream to obtain the Total HSA in the up- and downstream (y-axis). All samples were assayed for prion using a sandwich ELISA, and recording the absorbance values at 450nm (second y-axis).

25 Figure 2. Samples from the second phase of an IgG separation were taken from both up- and downstreams (U/S and D/S respectively) at 30 minute intervals. The samples were assayed for endotoxin using a LAL Chromogenic assay (Cape Cod Assoc.)

Figure 3. HSA was purified from endotoxin spiked plasma. Samples were taken from up - and downstream at 30 minute intervals during a 90 minute purification (x-axis). Analysis of the samples using a LAL Chromogenic assay was performed to establish the
30 endotoxin concentration (y-axis) in the samples.

Figure 4. Four to 25% native gel electrophoresis of samples from an HSA

purification from endotoxin spiked plasma. Lane 1 contains molecular weight markers. Lane 2 contains starting plasma sample, Lanes 3-5 contain upstream samples at time 30, 60, and 90 minutes. Lanes 6-9 contain downstream samples at time 0, 30, 60 and 90 minutes, respectively.

5 Modes for Carrying Out the Invention

Virus removal during plasma protein purification using Gradiflow technology

Contamination with virus is a major concern when purifying plasma proteins, such as IgG and human serum albumin (HSA). A contaminant virus can potentially infect a patient receiving the contaminated plasma products. A virus that infects bacteria is known as a phage, and they are readily detected by examining culture plates for cleared zones in a coating or lawn of bacteria.

Aim. To isolate IgG, HSA, and Fibrinogen from human plasma spiked with virus, using the Gradiflow, with simultaneous removal of the contaminating virus.

IgG purification procedure

IgG is the most abundant of the immunoglobulins, representing almost 70% of the total immunoglobulins in human serum. This class of immunoglobulins has a molecular mass of approximately 150kDa and consists of 4 subunits, two of which are light chains and two of which are heavy chains. The concentration of IgG in normal serum is approximately 10mg/ml.

IgGs are conventionally purified using Protein A affinity columns in combination with DEAE-cellulose or DEAE-Sephadex columns. The main biological contaminants in IgG isolations are β -lipoprotein and transferrin. The product of conventional protein purification protocols is concentrated using ultrafiltration. Immunoaffinity can also be used to isolate specific IgGs.

Method: Platelet free plasma was diluted one part in three with Tris-borate, pH 9.0 running buffer and placed in the upstream of Gradiflow and spiked with either Lambda or T7 phage to a concentration of $\sim 10^8$ pfu/ml (plaque forming units/ml). A potential of 250V was placed across a separating membrane with a molecular weight cut off of 200kDa (3kDa restriction membranes). A membrane of this size restricts IgG migration whilst allowing smaller molecular weight contaminants to pass through the membrane, leaving IgG and other large molecular weight proteins in the upstream. A second purification phase was carried out using a

GABA/Acetic acid buffer, pH 4.6 with a 500kDa cut off separating membrane (3kDa restriction membranes). A potential of 250V reversed polarity was placed across the system resulting in IgG migration through the membrane leaving other high molecular weight contaminants upstream.

5 Examination of samples taken at 30 minutes intervals was made on reduced SDS-PAGE 4-25% gels.

Virus testing

10 One hundred and fifty μ l taken at each time point sample was mixed with 100 μ l of appropriate *Escherichia coli* culture (Strain HB101 was used for T7 and strain JM101 for Llambda). The mixtures were incubated for 15 minutes at 37°C before each was added to 2.5 ml of freshly prepared molten soft agar, and vortexed. The mixtures were poured over culture plates of Luria Agar and incubated at 37°C overnight. The plates were inspected for the presence of virus colonies (plaques) in the lawn of *E. coli*. The number of plaques was recorded or if the virus had infected the entire *E. coli* population the result was recorded as confluent lysis.

HSA purification procedure

20 Albumin is the most abundant protein component (50mg/m.) in human plasma and functions to maintain blood volume and oncotic pressure. Albumin regulates the transport of protein, fatty acids, hormones and drugs in the body. Clinical uses include blood volume replacement during surgery, shock, serious burns and other medical emergencies. Albumin is 67kDa and has an isoelectric point of approximately 4.9. The protein consists of a single subunit and is globular in shape. About 440 metric tons of albumin is used annually internationally with worldwide sales of US \$1.5 billion. Albumin is currently purified using Cohn fractionation and commercial product contains many contaminants in addition to multimers of albumin. The high concentration, globular nature and solubility of albumin make it an ideal candidate for purification from plasma using Gradiflow technology.

30 *Method:* Pooled normal plasma was diluted one in three with Tris-Borate (TB) running buffer, pH 9.0 and spiked with $\sim 10^8$ pfu/ml of Llambda or T7 phage. The mixture was placed in the upstream of a Gradiflow apparatus. Albumin was isolated from platelet free plasma in a one-phase process using the charge of albumin at a pH above its isoelectric point (pI) and its

molecular weight. Thus, a cartridge with a 75kDa cutoff separation membrane was placed between two 50kDa cutoff restriction membranes. The albumin was removed from high molecular weight contaminants by its migration through the separation membrane whilst small molecular weight contaminants dissipated through the 50kDa restriction membrane. Samples were taken at regular intervals throughout a 90 minutes run.

The presence of the purified HSA in the downstream was demonstrated by examination by SDS-PAGE. Virus was detected as previously described above.

Fibrinogen purification procedure:

Commercially, fibrinogen has a role as fibrin glue, which is used to arrest bleeding and assist in the wound healing process. Fibrinogen is an elongated molecule of 340kDa that consists of three non-identical subunit pairs that are linked by a disulphide knot in a coiled coil conformation. The isoelectric point of fibrinogen is 5.5 and it is sparingly soluble when compared with other plasma proteins.

Fibrinogen is conventionally purified from plasma by a series of techniques including ethanol precipitation, affinity columns and traditional electrophoresis. This process takes about 48-72 hours and the harsh physical and chemical stresses placed on fibrinogen are believed to denature the molecule, resulting in activity that is removed from that of fibrinogen in plasma.

Cryo-precipitation is the first step in the production of factor VIII and involves the loss of most of the fibrinogen in plasma. Processing of this waste fibrinogen is of considerable interest to major plasma processors and provides an opportunity to demonstrate the rapid purification of fibrinogen from cryo-precipitate using the Gradiflow.

Method: Cryo-precipitate 1, produced by thawing frozen plasma at 4°C overnight was removed from plasma by centrifugation at 10000xg at 4°C for 5 minutes. The precipitate was re-dissolved in Tris-Borate buffer (pH 9.0) and placed in the upstream of a Gradiflow apparatus. The upstream was spiked with either Lambda or T7 phage to a concentration of $\sim 10^8$ pfu/ml. A potential of 250V was applied across a 300kDa cutoff cartridge and run for 2 hours. The downstream was replaced with fresh buffer at 30 minute intervals. A second phase was used to concentrate the fibrinogen through a 500kDa cutoff separation membrane at pH 9.0. The downstream was harvested at 60 minutes. The product was dialysed against PBS pH 7.2 and analysed for clotting activity by the addition of calcium and thrombin (final concentrations

10mM and 10NIH unit/ml respectively).

The presence of purified fibrinogen was confirmed by examination on reduced SDS PAGE 4-25% gels. The presence of either T7 or Llambda in the time point samples was tested using the previously described method.

5

Results of IgG, HSA and fibrinogen purification:

The procedures described successfully purified IgG, albumin and fibrinogen as judged by electrophoresis. Neither T7 nor Llambda phage were detected in the downstream products, but were present in the upstream samples.

10 **Prion removal during plasma protein purification using Gradiflow technology**

There is an international concern regarding the contamination of plasma proteins by prion protein. Prion is a glycoprotein of 27-33kDa in size which occurs naturally in many human derived materials, including white blood cells, platelets, plasma and plasma proteins preparations, e.g., HSA, IgG, FVIII and fibrinogen. Prion can become folded abnormally and cause neurological disorders such as Creutzfeld-Jacob disease (CJD) and Kuru. Currently, there is much concern regarding the transmission of these diseases via transfusion and plasma protein fractions administered clinically.

Aim: To isolate NSA from human plasma using the Gradiflow, with simultaneous removal of prion.

Method: Pooled platelet rich plasma was diluted one in two with Tris-Borate (TB) running buffer, pH 9.0 and was placed in the upstream of a Gradiflow apparatus. Albumin was isolated from platelet free plasma using the charge of albumin at a pH above its pI and its molecular weight. Thus, a cartridge with a 75kDa cutoff separation membrane was placed between two 50kDa cutoff restriction membranes. The albumin was removed from high
25 molecular weight contaminants by its migration through the separation membrane whilst small molecular weight contaminants dissipated through the 50kDa restriction membrane. Samples were taken at 20 minute intervals throughout a 240 minute run. The buffer stream and cartridge were replaced after the initial two hours, with identical solutions and cartridge.

30 The presence of the purified HSA in the downstream was demonstrated by examination by SDS-PAGE, and was measured using a Bromocresol Green Assay (purchase from Trace Scientific. Prion was tested for in both up - and down-stream samples using a sandwich ELISA

comprised of prion specific antibodies obtained from Prionics Inc (Switzerland).

Albumin quantitation

Fifty μl of each time point sample was diluted with 50 μl of PBS buffer. A 20 μl aliquot of each diluted sample was placed in a microplate well. A standard curve of the kit calibrator from a maximum concentration of 40mg/ml was prepared using PBS as the diluent. The standard curve dilutions were also placed in the microplate (2 μl plasma/well). The bromocresol green reagent was added to all the wells (200 μl /well) and the absorbance at 630nm was read using a Versamax microplate reader. The standard curve was drawn on a linear scale and the concentration of albumin in the up and downstream samples were read from the curve. The volume in the appropriate stream at the time of sampling was multiplied by the concentration of each sample. Thus providing a value for the total HSA present in each stream.

Prion detection

A solution of 5 $\mu\text{g}/\text{ml}$ monoclonal antibody denoted 6H4 (Prionics, Inc. Switzerland) in a 10mM carbonate buffer was added to the wells of a microplate (100 μl /well), and incubated overnight at 4°C. The antibody was later decanted and the wells washed three times with 250 μl /well of a PBS solution containing 0.1%(v/v) Tween 20. The plate wells were blocked by incubating at room temperature for 30 minutes with 200 μl /well of PBS/T20 containing 1% albumin. The plate was again washed three times with 250 μl /well of PBS/T20 before the up- and down-stream time point samples were added (100 μl /well). The samples were incubated for 1-2 hours at room temperature before being dispensed, and the plate washed three times as previously described. A solution of prion-specific polyclonal antibody, denoted R029 (Prionics Inc. Switzerland) was diluted at 1:1000(v:v) in PBS/T20, and added to the wells of the plate (100 μl /well). The mixture was incubated for 1-2 hours at room temperature, before being decanted. The plate was washed three times and 100 μl /well of horseradish peroxidase conjugated polyclonal anti-rabbit IgG antiserum (purchased from Dakopatts) was added. The conjugate was incubated for 30-60 minutes at room temperature and then removed. Any bound HRP conjugate was detected using o-tolidine substrate solution (100 μl /well) and the reaction stopped by addition of 3M HCl (50 μl /well). The developed colour was measured at 450nm in a Versamax plate reader.

Results

Albumin was transferred to the downstream and was detected in the BCG assay (Figure 1), and visualized on a native 8-16% electrophoresis gel. Decreasing quantities of Prion were detected in the upstream during the time-course, and no Prion was detected in the downstream samples.

Endotoxin removal during plasma protein purification using Gradiflow technology

Contamination with bacterial endotoxin is a major concern when purifying plasma proteins, such as IgG and HSA. Endotoxins are a lipopolysaccharide derived from the lipid membrane of gram negative bacteria. The presence of endotoxin in a human blood fraction therapeutic can lead to death of the receiving patients.

Aim: To isolate IgG and HSA from human plasma spiked with endotoxin, using the Gradiflow, with simultaneous removal of endotoxin.

IgG purification procedure

Method: Platelet free plasma was diluted one part in three with Tris-borate, pH 9.0 running buffer and placed in the upstream of a Gradiflow apparatus and spiked with purified *E. coli* endotoxin to a concentration of 55ng/ml. A potential of 250V was placed across a separating membrane with a molecular weight cut off of 200kDa (3kDa restriction membranes). A membrane of this size restricts IgG migration whilst allowing smaller molecular weight contaminants to pass through the membrane, leaving IgG and other large molecular weight proteins in the upstream. A second purification phase was carried out using a GABA/Acetic acid buffer, pH 4.6 with a 500kDa cut off separating membrane (3kDa restriction membranes). A potential of 250V reversed polarity was placed across the system resulting in IgG migration through the membrane leaving other high molecular weight contaminants upstream.

Examination of samples taken at 30 minutes intervals was made on reduced SDS-PAGE 4-25% gels. Endotoxin was tested for using a LAL Pyrochrome Chromogenic assay purchased from Cape Cod Associates. All samples were diluted 1 in 10 and the endotoxin assay was performed according to the manufacturer instructions.

HSA purification procedure

Method: Pooled normal plasma was diluted one in three with Tris-Borate (TB)

running buffer, pH 9.0 and spiked with 55ng/ml of purified endotoxin. The mixture was placed in the upstream of a Gradiflow apparatus. Albumin was isolated from platelet free plasma in a one-phase process using the charge of albumin at a pH above its pI and its molecular weight. Thus, a cartridge with a 75kDa cutoff separation membrane was placed between two 50kDa cutoff restriction membranes. The albumin was removed from high molecular weight contaminants by its migration through the separation membrane whilst small molecular weight contaminants dissipated through the 50kDa restriction membrane. Samples were taken at regular intervals throughout a 90 minute run.

The presence of the purified HSA in the downstream was demonstrated by examination by SDS-PAGE. Endotoxin was tested for in both up - and down-stream samples using a LAL Chromogenic assay supplied by Cape Cod Associates. All samples were diluted 1 in 10 and the endotoxin assay was performed according to the manufacturer instructions.

Results of IgG and HSA purification

Up and downstream samples taken at 30 minute intervals during the second phase of an IgG purification from endotoxin spiked plasma were tested for endotoxin using a LAL Chromogenic assay. The results showed that the endotoxin was almost entirely found in the upstream at all time points (Figure 2). The downstream contained only 0.7% of the initial endotoxin. Reduced SDS-PAGE examination showed that IgG had been successfully isolated in the downstream.

Analysis of samples taken at 30 minute intervals during the purification of HSA from plasma spiked with endotoxin found the majority of endotoxin remained in the upstream. Only 4% of the total endotoxin was found in the downstream at the end of the run (Figure 3). Native PAGE examination confirmed the presence of purified HSA in the downstream samples (Figure 4).

Bacteria removal during plasma protein purification using Gradiflow technology

Contamination with bacteria is a major concern when purifying plasma proteins, such as IgG and HSA. Contaminant bacteria can potentially infect a patient receiving the plasma products, or during pasteurisation of the products the bacteria dies releasing dangerous endotoxins, that are harmful to the patient. Bacteria are easily detected by culturing samples on nutrient agar plates.

Aim: To isolate IgG. and HSA, from human plasma spiked with bacteria, using the Gradiflow.

IgG purification procedure

5 *Method:* Platelet free plasma was diluted one part in three with Tris-borate, pH 9.0 running buffer and placed in the upstream of Gradiflow and spiked with *E. coil* to a concentration of 4×10^8 cells/ml. A potential of 250V was placed across a separating membrane with a molecular weight cut off of 200kDa (100kDa restriction membranes). A membrane of this size restricts IgG migration whilst allowing smaller molecular weight contaminants to pass
10 through the membrane, leaving IgG and other large molecular weight proteins in the upstream. A second purification phase was carried out using a GABA/Acetic acid buffer, pH 4.6 with a 500kDa cut off separating membrane (3kDa restriction membranes). A potential of 250V reversed polarity was placed across the system resulting in IgG migration through the membrane leaving other high molecular weight contaminants upstream.

Examination of samples taken at 30 minutes intervals was made on reduced SDS-PAGE 4-25% gels.

Bacteria testing

Twenty μ l of upstream or 100 μ l of downstream samples were spread plated onto Luria agar culture plates. The plates were incubated for 24 hours at 37°C, and the number of colonies was counted.

HSA purification procedure

Method: Pooled normal plasma was diluted one in three with Tris-Borate (TB) running buffer, pH 9.0 and spiked with $\sim 4 \times 10^8$ cells/ml of *E. coli*. The mixture was placed in the upstream of a Gradiflow apparatus. Albumin was isolated from platelet free plasma in a one-phase process using the charge of albumin at a pH above its pI and its molecular weight. Thus a cartridge with a 75kDa cutoff separation membrane was placed between two 50kDa cutoff restriction membranes. The albumin was removed from high molecular weight contaminants by its migration through the separation membrane whilst small molecular weight contaminants dissipated through the 50kDa restriction membrane. Samples were taken at regular intervals throughout a 90 minutes run.

The presence of the purified HSA in the downstream was demonstrated by examination by SDS-PAGE. Bacteria were detected as previously described above.

Results of IgG and HSA purification

The procedures described successfully purified IgG, and albumin as judged by electrophoretic examination. The downstream samples containing the purified protein products did not contain detectable *E. coli* colonies, while the upstream samples produced greatly in excess of 500 colonies/plate.

CONCLUSION

It is possible to purify proteins such as IgG, albumin and fibrinogen from plasma, while simultaneously removing contaminating virus by the methods according to the present invention.

Prion present in plasma can be moved across a 75kDa separation membrane with albumin, however, unlike albumin, the prion is not retained by the 50kDa restriction membrane. Thus, albumin can be purified from plasma with simultaneous removal of Prion protein.

Evidence has been provided by the present inventors that it is possible to purify proteins such as IgG and albumin from plasma. while simultaneously removing endotoxin contamination in the starting plasma using the Gradiflow technology.

Furthermore, it has been found that it is also possible to purify proteins such as IgG, and
5 albumin from plasma, while simultaneously removing contaminating bacteria.

It will be appreciated by persons skilled in the art that numerous variations and/or modifications may be made to the invention as shown in the specific embodiments without departing from the spirit or scope of the invention as broadly described. The present embodiments are, therefore, to be considered in all respects as illustrative and not restrictive.

10

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000